

64420

Trench Soil
331.1 grams

DRAFT



Figure 1: Trench at station 4, Apollo 16. AS16-107-17461.

CDR: Okay, Houston. I'm digging an exploratory trench right here to see if material is black. – the material is not white. It's just the same as it - I've gone down about shovel width, and it's all the same material, and I don't see any layering in it or anything.

CDR: I've got a sample out of the deepest part of this trench that I'm digging – bag 399

Introduction

Sample 64420 is from the bottom of a 25 cm deep trench (figure 1) at the bottom of a small double crater on the rim of a 15 meter crater (Cinco 'a') on Stone Mountain (figure 2). It included one small (14 g) fragment of 'dilithologic breccia' (64425).

Double drive tube 64002/1, a large reference soil 64500 and a rake sample (64535 - 64589) were collected nearby. Samples from station 4, high up on Stone Mountain, should primarily be materials from the Descartes Formation. However, it has not proven easy to distinguish Descartes material from that of Cayley. A complication is that this area is on a ray that extends




Figure 2: Plan view of samples collected at station 4 (highest point on Stone Mountain, Apollo 16).

Modal content of soil 64421.

From Butler et al. 1973.

74 – 53 micron

Olivine	0.6
Pyroxene	1.1
Plagioclase	13.6
Glass other	10.6
Rock fragments	22.9
Welded fragments	52.6




Figure 3 : Compositions of lunar soils with 64420 plotted.




Figure 4: Normalized rare-earth-element diagram for 64421 compared with other Apollo 16 soils.

back to South Ray Crater. It is thought that dimict breccias (e.g. 64435, 64475, 64476 as well as 64425) may be ejecta from South Ray Crater (James and Lindstrom 1991).

It has high maturity, especially considering that it's from the bottom of a trench.

Petrography

Morris (1978) reported a maturity index $Is/FeO = 83$ (figure 5) and the mode shows a high percentage of ‘welded fragments’ (agglutinates). Butler et al. (1973) and Graf (1993) report the grain size analysis based on sieving (figure 7).

Note that sample 64425 is a dimict breccia from this soil (James and Lindstrom 1991). Breccias 64535 and 64475 are large pieces of ‘dilithologic breccia’ from the same location.

Chemistry

The chemical composition of 64421 (table 1) is similar to that of soil sample 64501 and the drive tube 64002 – 1 (figures 3 and 4). Duncan et al. (1973) calculate that this soil is ~ 76 % anorthosite, 11 % KREEP, 9 % mare basalt and 3 % meteorite, while Kempa et al. (1980) calculate ~ 60 % anorthosite, 30 % low-K Fra Mauro (impact melt) and 10 % mare basalt (using components found in the sample itself).

Moore et al. (1973) reported relatively high total carbon for this sample (280 ppm), while Epstein and Taylor (1975) found more normal amounts (figure 6). Muller (1973) determined 124 ppm nitrogen, consistent with the Epstein and Taylor results.

Rhodes et al. (1975) found a systematic change in composition of agglutinates; apparently refuted by Hu and Taylor (1977).

Cosmogenic isotopes and exposure ages

Clark and Keith (1973) determined the cosmic-ray-induced activity for $^{26}\text{Al} = 111 \text{ dpm/kg}$, $^{22}\text{Na} = 39 \text{ dpm/kg}$ and $^{56}\text{Co} = 19 \text{ dpm/kg}$. Nishiizumi et al. (1983) determined ^{53}Mn along the length of the core 64002. They seem to have found that the top of the core had been recently “disturbed”, which would also apply to the trench soil (which seem to be unusually mature at depth).

Kirsten et al. (1973) reported a Ne exposure age of 210 m.y.

Other Studies

Behrmann et al. (1973) and Poupeau et al. (1975) found an abundance of cosmic ray tracks in this mature soil from the bottom of the trench (figure 7)! Bogard and Nyquist (1973), Kirsten et al. (1973) and Wieler et al. (1980) reported rare gases and their isotopes. Muller (1973) discusses the positive correlation of bound N with C, He and Ne contents – all derived from solar wind implantation.




Figure 5: C from Epstein and Taylor (1974); maturity from Morris (1978).

References 64420

Bansal B.M., Church S.E., Gast P.W., Hubbard N.J., Rhodes J.M. and Weismann H. (1972) The chemical composition of soil from the Apollo 16 and Luna 20 sites. Earth Planet. Sci. Lett. 17, 29-35.

Behrman C.J., Crozaz G., Drozd R., Hohenberg C., Ralston C., Walker R. and Yuhas D. (1973) Cosmic-ray exposure history of North Ray and South Ray material. Proc. 4th Lunar Sci. Conf. 1957-1974.

Bogard and Nyquist (1973) 40Ar/39Ar variations in Apollo 15 and 16 regolith. Proc. 4th Lunar Sci. Conf. 1975-1985.

Butler P. (1972) Lunar Sample Information Catalog Apollo 16. Lunar Receiving Laboratory. MSC 03210 Curator's Catalog, pp. 370.

Butler J.C., Greene G.M. and King E.A. (1973) Grain size frequency distribution and modal analysis of Apollo 16 fines. Proc. 4th Lunar Sci. Conf. 267-278.

Cirlin E.H. and Housley R.M. (1981) Distribution and evolution of Zn, Cd and Pb in Apollo 16 regolith samples




Figure 6: Carbon content from Epstein and Taylor (1974) and Moore et al. (1973) compared with other Apollo 16 samples.





Figure 7: Grain size distribution of 64421 (from Graf 1993; from data by Butler et al. 1973).

and the average U-Pb ages of the parent rocks. Proc. 12th Lunar Planet. Sci. Conf. 529-540.

Clark R.S. and Keith J.E. (1973) Determination of natural and cosmic ray induced radionuclides in Apollo 16 lunar samples. Proc. 4th Lunar Sci. Conf. 2105-2113.

Compston W., Vernon M.J., Chappell B.W. and Freeman R. (1973) Rb-Sr model ages and chemical composition of nine Apollo 16 soils (abs). Lunar Sci. IV, 158.

Table 1. Chemical composition of 64421.

reference weight	LSPET73	Wanke73	Bansal72	Muller75	Compston73	Rhodes75	Clark 73	Hubbard73
SiO ₂ %	44.88	(a) 45.14	(a)		45.01	(a) 44.97	(a)	44.68 (a)
TiO ₂	0.55	(a) 0.53	(a)	0.52 (d)	0.56	(a) 0.53	(a)	0.55 (a)
Al ₂ O ₃	27.6	(a) 27.8	(a)	27.4	(e) 27.62	(a) 27.82	(a)	27.63 (a)
FeO	5.03	(a) 4.95	(a)	4.89	(e) 5	(a) 4.71	(a)	4.9 (a)
MnO	0.06	(a) 0.06	(a)		0.07	(a) 0.05	(a)	0.08 (a)
MgO	5.35	(a) 5.24	(a)	5.39	(e) 5.47	(a) 5.36	(a)	5.33 (a)
CaO	15.81	(a) 15.81	(a)	15.7	(e) 16.01	(a) 15.71	(a)	15.69 (a)
Na ₂ O	0.39	(a) 0.44	(a)	0.46 (d)	0.45 (e)	0.44 (a)	0.58 (a)	0.47 (a)
K ₂ O	0.1	(a) 0.1	(a)	0.12 (d)	0.13 (b)	0.11 (a)	0.11 (a)	0.11 (a)
P ₂ O ₅	0.13	(a)			0.12	(a) 0.09	(a)	0.11 (a)
S %	0.07	(a)			0.08	(a) 0.05	(a)	0.07 (a)
<i>sum</i>								
Sc ppm		7.9	(b)			8.21	(b)	
V								
Cr	710	(a) 620	(b)	700 (d)		530	(b)	
Co		25	(b)			16.6	(b)	
Ni	316	(a) 330	(b)			210	(b)	
Cu								
Zn								
Ga								
Ge ppb								
As								
Se								
Rb	2.9	(a)		2.72 (d)	2.7 (b)	2.7		
Sr	172	(a) 160	(b)	173 (d)	177 (b)	165		
Y	42	(a) 40	(b)					
Zr	183	(a) 174	(b)	185 (d)				
Nb	11	(a) 9.5	(b)					
Mo								
Ru								
Rh								
Pd ppb								
Ag ppb								
Cd ppb								
In ppb								
Sn ppb								
Sb ppb								
Te ppb								
Cs ppm				0.13 (b)				
Ba	115	(b) 143	(d)	159 (b)				
La	12.3	(b) 12.5	(d)	13.4 (b)		10.1 (b)		
Ce	35	(b) 32.1	(d)			26 (b)		
Pr								
Nd		19.8 (d)						
Sm	5.7	(b) 5.61	(d)			4.7 (b)		
Eu	1.14	(b) 1.18	(d)			1.23 (b)		
Gd		6.67 (d)						
Tb	1.1	(b)				1.1 (b)		
Dy	6.8	(b) 7.5	(d)					
Ho	1.6	(b)						
Er		4.57 (d)						
Tm								
Yb	4	(b) 4.14	(d)			3.6 (b)		
Lu	0.57	(b) 0.61	(d)			0.49 (b)		
Hf	4	(b)				3.9 (b)		
Ta	0.5	(b)				0.4 (b)		
W ppb								
Re ppb								
Os ppb								
Ir ppb		9						
Pt ppb								
Au ppb		7.5						
Th ppm	2.8	(a)				1.6 (b)	2 (f)	
U ppm			0.66 (d)	0.58 (b)			0.62 (f)	

technique: (a) XRF, (b) INAA, (c) RNAA, (d) IDMS, (e) AA, (f) radiation counting

DesMarias D.J., Hayes J.M. and Meinschein W.G. (1973) The distribution in lunar soil of carbon released by pyrolysis. Proc. 4th Lunar Sci. Conf. 1543-1558.

DesMarias D.J., Hayes J.M. and Meinschein W.G. (1974) The distribution in lunar soil of hydrogen released by pyrolysis. Proc. 5th Lunar Sci. Conf. 1811-1822.

Duncan A.R., Erlank A.J., Willis J.P. and Ahrens L.H. (1973) Composition and inter-relationships of some Apollo 16 samples. Proc. 4th Lunar Sci. Conf. 1097-1113.

Epstein S. and Taylor H.P. (1973) The isotopic composition and concentration of water, hydrogen, and carbon in some Apollo 15 and 16 soils and in the Apollo 17 orange soil. Proc. 4th Lunar Sci. Conf. 1559-1575.

Epstein S. and Taylor H.P. (1975) Investigation of carbon, hydrogen, oxygen and silicon isotope and concentration relationships on the grain surfaces of a variety of lunar soils and in some Apollo 15 and 16 core samples. Proc. 6th Lunar Sci. Conf. 1171-1798.

Fruland R.M. (1983) Regolith Breccia Workbook. Curatorial Branch Publication # 66. JSC 19045.

Graf J.C. (1993) Lunar Soils Grain Size Catalog. JSC

Hubbard N.J., Rhodes J.M., Gast P.W., Bansal B.M., Shih C.-Y., Wiesmann H. and Nyquist L.E. (1973b) Lunar rock types: The role of plagioclase in non-mare and highland rock types. Proc. 4th Lunar Sci. Conf. 1297-1312.





Figure 7: Track density for samples affected by South Ray Crater (Behrman et al. 1973).

Hu H-N. and Taylor L-A. (1977) Lack of chemical fractionation in major and minor elements during agglutinate formation. Proc. 8th Lunar Sci. Conf. 3645-3656.

James O.B. (1981) Petrologic and age relations of the Apollo 16 rocks: Implications for subsurface geology and the age of the Nectaris Basin. Proc. 12th Lunar Planet. Sci. Conf. 209-233.



- James O.B. and Lindstrom M.M. (1991) Apollo 16 dimict breccias: I. Sample 64425 (abs). *Lunar Planet. Sci.* VII, 635-636.
- Janghorbani M., Miller M.D., Ma M-S., Chyi L.L. and Ehmann W.D. (1973) Oxygen and other elemental abundance data for Apollo 14, 15, 16 and 17 samples. *Proc. 4th Lunar Sci. Conf.* 1115-1126.
- Kempa M.J., Papike J.J. and White C. (1980) The Apollo 16 regolith: A petrographically-constrained chemical mixing model. *Proc. 11th Lunar Planet. Sci. Conf.* 1341-1355.
- Kerridge J.F., Kaplan I.R. and Petrowski C. (1975) Nitrogen in the lunar regolith: Solar origin and effects (abs). *Lunar Sci.* VI, 469-471.
- Kerridge J.F., Kaplan I.R. and Petrowski C. (1975) Evidence for meteoritic sulfur in lunar regolith. *Proc. 6th Lunar Sci. Conf.* 2151-2162.
- Kirsten T., Horn P. and Kiko J. (1973) $^{39}\text{Ar}/^{40}\text{Ar}$ dating and rare gas analysis of Apollo 16 rocks and soils. *Proc. 4th Lunar Sci. Conf.* 1757-1784.
- Korotev R.L (1981) Compositional trends in Apollo 16 soils. *Proc. 12th Lunar Sci. Conf.* 577-605.
- LSPET (1973) The Apollo 16 lunar samples: Petrographic and chemical description. *Science* 179, 23-34.
- LSPET (1972) Preliminary examination of lunar samples. Apollo 16 Preliminary Science Report. NASA SP-315, 7-1—7-58.
- Marvin U.B. (1972) Apollo 16 coarse fines (4-10 mm): Sample classification, description and inventory. *JSC Catalog*.
- Miller M.D., Pacer R.A., Ma M.-S., Hawke B.R., Lookhart G.L. and Ehmann W.D. (1974) Compositional studies of the lunar regolith at the Apollo 17 site. *Proc. 5th Lunar Sci. Conf.* 1079-1086.
- Moore C.B., Lewis C.F. and Gibson E.K. (1973) Total carbon contents of Apollo 15 and 16 lunar samples. *Proc. 4th Lunar Sci. Conf.* 1613-1623.
- Morris R.V., Score R., Dardano C. and Heiken G. (1983) Handbook of Lunar Soils. Two Parts. JSC 19069. Curator's Office, Houston
- Morris R.V. (1976) Surface exposure indicies of lunar soils: A comparative FMR study. *Proc. 7th Lunar Sci. Conf.* 315-335.
- Morris R.V. (1978) The surface exposure (maturity) of lunar soils: Some concepts and Is/FeO compilation. *Proc. 9th Lunar Sci. Conf.* 2287-2297.
- Muller O. (1973) Chemically bound nitrogen contents of Apollo 16 and Apollo 15 lunar fines. *Proc. 4th Lunar Sci. Conf.* 1625-1634.
- Muller O. (1975) Lithophile trace and major elements in Apollo 16 and 17 lunar samples. *Proc. 6th Lunar Sci. Conf.* 1303-1312.
- Poupeau G., Walker R.M., Zinner E. and Morrison D.A. (1975) Surface exposure history of individual crystals in the lunar regolith. *Proc. 6th Lunar Sci. Conf.* 3433-3448.
- Rhodes J.M., Adams J.B., Blanchard D.P., Charette M.P., Rodgers K.V., Jacobs J.W., Brannon J.C. and Haskin L.A. (1975) Chemistry of agglutinate fractions in lunar soils. *Proc. 6th Lunar Sci. Conf.* 2291-2308.
- Sutton R.L. (1981) Documentation of Apollo 16 samples. In USGS Prof. Paper 1048 (Ulrich et al. eds.)
- Taylor H.P. and Epstein S. (1973) O₁₈/O₁₆ and Si₃₀/Si₂₈ studies of some Apollo 15, 16 and 17 samples. *Proc. 4th Lunar Sci. Conf.* 1657-1679.
- Wieler R., Etique Ph., Signer P. and Poupeau G. (1980) Record of the solar corpuscular radiation in minerals from lunar soils: A comparative study of noble gases and tracks. *Proc. 11th Lunar Planet. Sci. Conf.* 1369-1393.
- Wänke H., Baddehausen H., Dreibus G., Jagoutz E., Kruse H., Palme H., Spettel B. and Teschke F. (1973) Multielement analysis of Apollo 15, 16 and 17 samples and the bulk composition of the moon. *Proc. 4th Lunar Sci. Conf.* 1461-1481.